

# Statistical analysis of the sampling design: FishPi case study on the biological sampling of the European hake fishery



Jose Castro, Nuno Prista, Lucia Zarauz, Jose Rodriguez, Manuela Azevedo, Laurent Dubroca, Alastair Pout, Joao Pereira, Chun Chen, Hans Gerritsen, Jon Elson, Ana Ribeiro, Patrik Börjesson, Kirsten Birch Håkansson, Sofie Nimmegeers




This work package was tasked with running **simulation models of the sampling** to test regional sampling designs for the fisheries exploiting regional stocks.


This simulation studies were prepared to **mimic the random selection of fishing trips** from the regional population of trips for that fishery.

The study focus on the statistical optimization of **landings estimates**.


Landings (mean 2013-2014) by vessel flag



Landings (mean 2013-2014) by sampling Institute



Field **"onShoreSampLoc"** indicates the **country responsible for the sampling** and is fundamental to organize a regional sampling program.



## CS4-total: list of scenarios

2-stage random sampling  
PSU: market/port and day  
SSU: trip

1. Baseline scenario: SRS
2. Scenario COUNTRY
3. Scenario INSTITUTE
4. Scenario REGIONAL stratified by PORT
5. Scenario COUNTRY and PORT
6. Scenario INSTITUTE and PORT

## CS4-southern: list of scenarios

2-stage random sampling  
PSU: market/port and day  
SSU: trip

1. Base line scenario: SRS
2. Scenarios REGIONAL.
  - 2a REGIONAL stratified by port
  - 2b REGIONAL stratified by quarter
  - 2c REGIONAL stratified by port-quarter
3. Scenarios COUNTRY
  - 3a COUNTRY stratified by port
  - 3b COUNTRY stratified by quarter
  - 3c COUNTRY stratified by port-quarter

**2013**

to set stratification  
and sample allocation

**2014**

to run simulations  
of each scenario

## Main criteria used in selecting the "best" scenarios

- Relative Bias:

Ratio between the estimated values (tons of hake) and the true value registered in 2014

- Relative Precision:

Ratio between the standard error (SE) of the estimated values of each scenario and the SE of Scenario 1 (*i.e.* 2-stage SRS not stratified)

- Coverage by country:

n PSUs, n SSUs, n SSUs with hake, tons hake available for sampling

- Coverage by domain:


n PSUs, n SSUs, n SSUs with hake, tons hake available for sampling

- Country
- Port and Quarter
- Stock
- FleetNat
- FoCatEu6

## CS4 - TOTAL SIMULATIONS RESULTS

Highest precision { scenario 4: **REGIONAL STRATIFIED BY PORT**  
scenario 5: **COUNTRY AND PORT**

Scenario	Sampling design	Precision
1	2-stage SRS (not stratified)	baseline
2	Stratified by Country	1.14
3	Stratified by Institute	1.44
4	REGIONAL stratified by PORT	<b>0.63</b>
5	Stratified by COUNTRY and PORT	<b>0.67</b>
6	Stratified by INSTITUTE and PORT	1.00



Institute	Current data		Port	Country/Port
	Nsamp_marketdays		SampTrips	SampTrips
ES_IEO	765	→	1035	652
PT	700	→	109	700
FR	597	→	882	597
GB_CEFAS	431	→	38	132
DK	113		104	113
IE	82	→	226	82
ES_AZTI	43	→	230	156
NL	38		3	38
GB_MS	11	→	189	344
SE	3		1	3


## CS4 - SOUTHERN SIMULATION RESULTS

Highest precision { scenario 2a: REGIONAL stratified by PORT  
scenario 2c: REGIONAL stratified by PORT-QUARTER

*The most precise overall but provide less coverage to some metiers and national fisheries.*


Scenario	sampling design	Precision
1	2-stage SRS WR (no stratified)	Baseline
2a	Regional stratified by port	<b>0.67</b>
2b	Regional stratified by quarter	1.05
2c	Regional stratified by port*quarter	<b>0.68</b>
3	Stratified by country	1.11
3a	Stratified by country and port	0.80
3b	Stratified by country and quarter	1.17
3c	Stratified by country and port*quarter	0.76
3o	Stratified by country (proportional to landings)	1.09

The second best-performing results were provided by scenarios 3a and 3c where the COUNTRY stratification is extended to PORT and PORT-QUARTER stratifications.




*Less precision in the estimation of total hake landings **but improved "tonnes of hake sampled" and "trips with hake sampled"** while providing **better coverage in some areas** with smaller hake landings.*

## CS4 - SOUTHERN SIMULATION RESULTS (examples: tonnes of Hke “sampled”)






## CS4 - SOUTHERN SIMULATION RESULTS (examples: market\_days without hake)



## CS4 - SOUTHERN SIMULATION RESULTS (examples: trips without hake)




## CS4-southern hake: Methodological insights

### Comparison between scenarios with all days and positive trips

#### Insights from CS4:

- Do we really know exactly in which market\*days and trips our target stocks will occur?
- Does this affect our perceptions on the quality of designs and how much Regional Sampling Plans can deliver?



## CS4-southern hake: Methodological insights

### Comparison between scenarios with all days and positive trips

#### Implications on size of the simulation dataset (2014)

	all market*days all_trips	all market*days pos_trips	pos market*days pos_trips
No. Markets	186	186	151
Market*days with hake landings	34849	34849	19074
No. Market*days	67890	67890	19074
No. Trips	584012	152465	103649

We ran simulations of these 3 setups using the same sampling effort  
and design Port2a (Regional Port Strata)

## CS4-southern hake: Methodological insights

### Comparison between scenarios with all days and positive trips

type	Expected value (mean) from 1000 sims			InterQuartile Range		
	all_marketdays all_trips	all_marketdays pos_trips	pos_marketdays pos_trips	all_marketdays all_trips	all_marketdays pos_trips	pos_marketdays pos_trips
tot_study	9592	9638	9606	1553	1068	641
ton_hke_in_ssu	207	503	769	30	46	53
n_ssu_with_hke_27.8.c	490	1283	2137	30	40	33
n_ssu_with_hke_27.9.a	472	1531	2687	29	44	33
tons_hke_sampled_27.8.c	160	353	542	29	43	49
tons_hke_sampled_27.9.a	47	150	227	8	15	17
n_ssu_with_hke_OTB_DEF_>=55_0_0	154	473	724	18	32	37
tons_hke_sampled_OTB_DEF_>=55_0_0	29	90	131	9	15	20

→ Sharp increase in expectations

## Considerations for selections of regional design

**Datasets to use:** define strata and sampling intensity and simulate on different datasets; consider thoroughly if datasets with only positive trips are applicable

**Precision:** analyze the scenarios with acceptable precision.

**Coverage by country:** analyze the **deviations that occur with respect to the current coverage.**

**Coverage by domain:** analyze the **deviations that occur in estimates of domains currently used by the National institutes to provide commercial data to ICES:** technical (métier), spatial (ICES Division), and time (quarter) disaggregation.

**Indicators of efficiency:** consider indicators like mean and variance of  $n\_ssu\_with\_spp$  and  $ton\_spp\_in\_sampled\_ssu$  per species, strata and post\_strata) should be a part of outputs so we approximate better what we can expect from design.

**Match real-life fleet operation conditions:** E.g., all days of the year should be included unless specific days are known not to have any landings; all fleets should be included unless specific fleets are known to have negligible landings with similar fish size/age structures.

**Adequacy to multiple end-user needs:** A regional sampling design should aim at **estimating accurate length/age composition** of the landings of the hake stocks for stock assessment purposes while at the same time providing **accurate estimates for many other stocks and fisheries subject to assessment.** The current study focused on estimating total landings of hake which are hardly a proxy for size/age.